

## A deterministic approach to the Skorokhod problem

by

Piernicola Bettiol

SISSA/ISAS

via Beirut, 2-4 - 34013 Trieste, Italy

e-mail: bettiol@sissa.it

**Abstract:** We prove an existence and uniqueness result for the solutions to the Skorokhod problem on uniformly prox-regular sets through a deterministic approach. This result can be applied in order to investigate some regularity properties of the value function for differential games with reflection on the boundary.

**Keywords:** Skorokhod problem, reflection on the boundary, differential inclusions, differential games, value function.

### 1. Introduction

The main purpose of the present article is to study the Skorokhod problem from a deterministic point of view and to get some properties on the value function for (zero-sum) differential games with reflection on the boundary.

Given  $T > 0$  and the initial data  $x \in K$ , by solving the deterministic Skorokhod problem, we mean finding the solution to the following differential inclusion:

$$\begin{cases} \dot{y}(t) \in f(t, y(t)) - \partial_- \psi_K(y(t)) \\ y(t) \in K \quad \forall t \in [0, T] \\ y(0) = x \in K \end{cases} \quad (1)$$

where  $\partial_- \psi_K$  is the subdifferential of the indicator function of  $K$ ,  $\psi_K$ .

More properly, the Skorokhod problem arises in the stochastic framework. It was studied and solved by Lions and Sznitman in Lions (1985) and Lions, and Sznitman (1984) for smooth domains (see also the reference therein for a more complete literature scene). See also Frankowska (1985), where a viability approach was used.

In the deterministic frame the analogous problem is often called “reflecting boundary problem”. Nevertheless, we prefer to keep the terminology used in Ishii (1988): the deterministic Skorokhod problem or (with a little abuse of language) simply the Skorokhod problem.

We assume that  $K \subset \mathbb{R}^n$  is a uniformly prox-regular set. This notion is equivalent to a very nice property: the projection to  $K$  is single-valued on the suitable (“tubular”) neighborhood of  $K$ .

For a rather complete description of such sets we refer the reader to Poliquin, Rockafellar, Thibault (2000) and the reference therein (see also Clarke, Stern, Wolenski, 1995). Since we deal with finite dimensional spaces, the notion of uniformly prox-normal sets coincides with the notion of Federer’s sets of positive reach (for properties of these sets see Federer, 1959; Delfour, Zolesio, 1994, 2004).

Examples of uniformly prox-regular sets are provided by  $K \subset \mathbb{R}^n$  such that  $\partial K \in \mathcal{C}^{1,1}$  or  $K$  convex.

Notice that system (1) without the term  $\partial_- \psi_K(y(t))$  becomes an usual Cauchy problem that admits a unique global Carathéodory solution, and this is the case when  $y(t) \in \text{Int}(K)$ . On the other hand, when  $y(t) \in \partial K$ , the trajectory  $y(\cdot)$  is pushed back along a direction that belongs to  $\partial_- \psi_K(y(t))$ , yielding a sort of reflection on the boundary. In fact, we can clarify better this concept by showing that system (1) really describes a reflection on the boundary of  $K$ . Indeed, for uniformly prox-regular sets the normal cone of  $K$  and the subdifferential of its indicator function coincide. Therefore, system (1) is equivalent to the viability problem

$$\begin{cases} \dot{y}(t) \in f(t, y(t)) - N_K(y(t)) \\ y(t) \in K \quad \forall t \in [0, T] \\ y(0) = x \in K \end{cases} \quad (2)$$

that, in turn, has the same set of solutions as the following one

$$\begin{cases} \dot{y}(t) = \Pi_{T_K(y(t))}(f(t, y(t))) \\ y(t) \in K \quad \forall t \in [0, T] \\ y(0) = x \in K. \end{cases} \quad (3)$$

The equivalence between (2) and (3) expresses quite well the reflecting role of the term  $\partial_- \psi_K(y(t))$  in system (1) (or of the term  $N_K(y(t))$  in (2)). Equivalence and existence results for these systems have been successfully studied in several papers and in different contexts: see for instance Henry (1973) who treats the case when  $K$  is convex and Cornet (1983) who assumes a tangential regularity of  $K$ .

More recently, in the frame of sweeping process Thibault (2003) provides existence results of viable solutions to (2) for  $K$  closed and with a reflection obtained by the Clarke cone. Furthermore, Serea (2003) gives an equivalence result dealing with a Mayer problem for controlled systems with reflection on the boundary of closed subsets  $K$  of  $\mathbb{R}^n$ ; she formulates system (2) by using the strict normal cone  $\hat{N}_K(y(t))$  providing an existence result for bounded sleek subsets and, moreover, also uniqueness when  $K$  is bounded and proximal retract.

In the present paper after introducing some preliminaries in Section 2, we devote Section 3 to a proof of an existence and uniqueness result using an adapted version of an approach due to Ishii in (1988): this approach yields not only a constructive proof of the trajectory that solves the deterministic Skorokhod problem (1), but also allows us to get existence and uniqueness also for not necessarily bounded prox-regular sets. Actually, one can obtain existence and uniqueness of the solution to our problem by applying different (maybe easier) approaches: for instance, by facing system (3) and by using an approach due to Viability Theory (see Remark 4). On the other hand, here our aim is to provide the unique solution to (1) by investigating directly the differential inclusion (1) and without invoking existence theorems for equivalent problems.

Finally, in the last section, we investigate the differential games with reflection on the boundary: once we fix the the controls of the players, the controlled Skorokhod problem becomes a system like (1). So, we show that the Lipschitz dependence of the solutions to the Skorokhod problem on the initial data provides a Hölder regularity for the value function. We also mention how this regularity could be useful to study the limiting behaviour of the value function, namely the so-called ergodic problem (see, for instance, Bardi and Capuzzo-Dolcetta, 1997).

## 2. Basic notions and assumptions

Let  $\mathbb{R}^n$  be endowed with the euclidean norm  $|\cdot|$ . Given  $K \subset \mathbb{R}^n$ , we denote the distance function from a point  $x \in \mathbb{R}^n$  to  $K$  by  $d_K(x) := \inf_{y \in K} |y - x|$ . We recall that the projection on  $K$ ,  $\Pi_K : \mathbb{R}^n \rightarrow K$ , is defined by

$$\Pi_K(x) := \{y \in K : |x - y| = d_K(x)\}.$$

We denote the closed ball of radius  $r$  centered in  $x_0 \in \mathbb{R}^n$  by  $B(x_0; r) := \{x \in \mathbb{R}^n : |x_0 - x| \leq r\}$ . For the closed unit ball in  $\mathbb{R}^n$  we simply write  $B$ . A set  $C \subset \mathbb{R}^n$  is called a cone if it is nonempty and for all  $\lambda \geq 0$  and  $v \in C$  we have  $\lambda v \in C$ . The negative polar cone of  $C$  is defined by

$$C^- = \{\xi \in \mathbb{R}^n \mid \langle \xi, v \rangle \leq 0, \quad \forall v \in C\}$$

where  $\langle \cdot, \cdot \rangle : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  is the canonical scalar product in  $\mathbb{R}^n$ .

**DEFINITION 1** *Let  $K$  be a subset of  $\mathbb{R}^n$  and  $x \in \overline{K}$ . Then*

1. *the contingent cone to  $K$  at  $x$ ,  $T_K(x)$ , is defined by*

$$T_K(x) := \left\{ v \mid \liminf_{h \rightarrow 0^+} \frac{d_K(x + hv)}{h} = 0 \right\};$$

2. *the proximal normal cone to  $K$  at  $x$ ,  $N_K^P(x)$ , is defined by*

$$N_K^P(x) := \{ \xi \mid \exists M > 0 \text{ such that } \langle \xi, y - x \rangle \leq M|y - x|^2 \quad \forall y \in K \};$$

3. the limiting normal cone (also called general cone of normals) to  $K$  at  $x$ ,  $N_K(x)$ , is defined by

$$N_K(x) := \left\{ \xi : \exists x_n \rightarrow_K x \text{ and } \xi_n \rightarrow \xi \text{ with } \xi_n \in N_K^P(x_n) \right\}.$$

For many properties on tangent and normal cones we refer the reader for instance to Aubin, Frankowska (1990), Rockafellar, Wets (1998) and Vinter (2000).

In general we have  $\{0\} \subset N_K^P(x) \subset N_K(x) \subset T_K^-(x)$ ;  $N_K^P(x)$  is convex,  $N_K(x)$  is closed and  $T_K^-(x) = \overline{\text{co}}N_K(x)$  (if  $Y \subset \mathbb{R}^n$ , then  $\overline{\text{co}}Y$  denotes the closed convex hull of  $Y$ ). We recall also that a closed set  $K \subset \mathbb{R}^n$  is called *sleek* if the set-valued map

$$K \ni x \rightsquigarrow T_K(x)$$

is lower semicontinuous.

In the present paper for the Skorokhod problem (1) we consider quite regular sets, that turn out to be also sleek (see Proposition 1 below): indeed, we assume that for some  $r_0 > 0$  fixed the closed set  $K \subset \mathbb{R}^n$  is *uniformly prox-regular* with constant  $\frac{1}{r}$  for any  $r \in (0, r_0)$  (or also called proximal  $r_0$ -retract set), namely, whenever  $x \in K$ ,  $r \in (0, r_0)$  and  $\xi \in N_K(x)$  with  $|\xi| < 1$ , then  $x$  is the unique nearest point of  $K$  to  $x + \frac{1}{r}\xi$  (see for instance Poliquin, Rockafellar, Thibault, 2000).

Prox-regular sets are characterized in several interesting ways and have a lot of important properties. In the following Proposition we recall some of them referring the reader to the article Poliquin, Rockafellar, Thibault (2000) for the proofs (in particular see Theorem 4.1 and Corollary 4.6).

**PROPOSITION 1** (Poliquin, Rockafellar, Thibault, 2000) *Let  $K$  be a closed subset of  $\mathbb{R}^n$  and  $r_0 > 0$ . Then, the following properties are equivalent:*

1.  $K$  is uniformly prox-regular with constant  $\frac{1}{r}$  for any  $r \in (0, r_0)$ ;
2.  $\Pi_K$  is single-valued on the “tubular” neighborhood of  $K$ ,  $U_K(r_0) := \{x \in \mathbb{R}^n : 0 < d_K(x) < r_0\}$ ;
3. every nonzero proximal normal to  $K$  at any point  $x \in K$  can be realized by an  $r_0$ -ball, namely for any  $x \in K$  and  $0 \neq \xi \in N_K^P(x)$  one has

$$K \cap \text{Int} \left\{ x + r_0 \left( \frac{\xi}{|\xi|} + B \right) \right\} = \emptyset;$$

4. whenever  $x_i \in K$  and  $\xi_i \in N_K^{r_0}(x_i) = N_K(x_i) \cap \text{Int}(B(0, r_0))$  we have

$$\langle \xi_1 - \xi_2, x_1 - x_2 \rangle \geq -|x_1 - x_2|^2.$$

Moreover, if one of the above equivalent properties occurs, then for all  $x \in K$

$$N_K^P(x) = N_K(x) = T_K^-(x).$$

REMARK 1 *Property 3 of Proposition 1 can be rephrased in the useful following way: given  $x \in K$ ,  $0 \neq \xi \in N_K^P(x)$  if and only if*

$$\frac{1}{2r_0}|y - x|^2 \geq \left\langle \frac{\xi}{|\xi|}, y - x \right\rangle \quad \forall y \in K.$$

So, for all  $x \in K$ ,  $\xi \in N_K^P(x)$  we have

$$\xi \cdot (x - y) + C_0|\xi||y - x|^2 \geq 0 \quad \forall y \in K \tag{4}$$

where  $C_0 = \frac{1}{2r_0}$ . An immediate consequence of (4) is that for any  $x_1, x_2 \in K$ ,  $\xi_1 \in N_K^P(x_1)$  and  $\xi_2 \in N_K^P(x_2)$

$$\langle \xi_1 - \xi_2, x - y \rangle + C_0(|\xi_1| + |\xi_2|)|x_1 - x_2|^2 \geq 0. \tag{5}$$

REMARK 2 *Proposition 1 yields that  $K$  is sleek. Property 4 of Proposition 1 says that the set-valued map  $K \ni x \rightsquigarrow N_K^{r_0}(x) + x$  is monotone (i.e. if  $\zeta_i \in N_K^{r_0}(x_i) + x_i$  for  $i = 1, 2$ , then  $\langle \zeta_1 - \zeta_2, x_1 - x_2 \rangle \geq 0$ ).*

Thanks to Proposition 1, henceforth, we simply talk about normal cones  $N_K(x)$  without specifying the type, since all the introduced notions coincide.

By  $\psi_K : \mathbb{R}^n \rightarrow [0, +\infty]$  we denote the indicator function of  $K$ , namely

$$\psi_K(x) = \begin{cases} 0 & \text{if } x \in K \\ +\infty & \text{if } x \notin K. \end{cases}$$

Note that  $\psi_K$  is lower semicontinuous on  $\mathbb{R}^n$  because  $K$  is closed.

Now, we introduce the notion of (local) subdifferential.

DEFINITION 2 *Given a function  $\psi : \mathbb{R}^n \rightarrow \overline{\mathbb{R}} := \mathbb{R} \cup \{\pm\infty\}$  and a point  $x_0$  such that  $\psi(x_0) \neq \pm\infty$ , the (local) subdifferential of  $\psi$  at a point  $x_0$  is so defined*

$$\partial_- \psi(x_0) := \left\{ \xi \in \mathbb{R}^n : \liminf_{x \rightarrow x_0} \frac{\psi(x) - \psi(x_0) - \langle \xi, x - x_0 \rangle}{|x - x_0|} \geq 0 \right\}.$$

We recall that by solving the deterministic Skorokhod problem we mean to find a solution to the following differential inclusion:

$$\begin{cases} \dot{y}(t) \in f(t, y(t)) - \partial_- \psi_K(y(t)) \\ y(t) \in K \quad \forall t \in [0, T] \\ y(0) = x \in K. \end{cases} \tag{6}$$

In the present paper  $f$  is a function so defined

$$\begin{aligned} f : \mathbb{R} \times \mathbb{R}^n &\longrightarrow \mathbb{R}^n \\ (t, x) &\longmapsto f(t, x) \end{aligned}$$

and we assume that

$$\left\{ \begin{array}{l} (i) \quad f(t, \cdot) \text{ is continuous for all } t, \\ (ii) \quad f \text{ is (Lebesgue) measurable with respect to } t, \\ (iii) \quad |f| < M \text{ for some } M > 0; \\ (iv) \quad \exists L > 0 \text{ such that} \\ \quad \langle f(t, x_1) - f(t, x_2), x_1 - x_2 \rangle \leq L|x_1 - x_2|^2 \quad \forall x_1, x_2 \in \mathbb{R}^n, t \in \mathbb{R}. \end{array} \right. \quad (7)$$

The regularity (in the Clarke sense) of the set  $K$  provides also a sort of regularity of the indicator function  $\psi_K$ , obtaining the following lemma (for the details of the proof, see for instance Rockafellar, Wets, 1998, Chapter 8, section B).

LEMMA 1 *If  $K \subset \mathbb{R}^n$  is uniformly prox-regular with constant  $\frac{1}{r}$  for any  $r \in (0, r_0)$  then we have*

$$\partial_- \psi_K(x) = N_K(x) \quad \forall x \in K.$$

Thus, thanks to Lemma 1, we can rewrite system (6) by means of the (viability) differential inclusion

$$\left\{ \begin{array}{l} \dot{y}(t) \in f(t, y(t)) - N_K(y(t)) \\ y(t) \in K \quad \forall t \in [0, T] \\ y(0) = x \in K. \end{array} \right. \quad (8)$$

The set of solutions of system (8) is the same as the following one (see, for instance, Serea, 2003, for the proof)

$$\left\{ \begin{array}{l} \dot{y}(t) = \Pi_{T_K(y(t))}(f(t, y(t))) \\ y(t) \in K \quad \forall t \in [0, T] \\ y(0) = x \in K. \end{array} \right. \quad (9)$$

Let us recall the definition of the inf-convolution of the function  $\psi_K(x)$  (see, for instance, Bardi, Capuzzo-Dolcetta, 1997, for this notion): for all  $\varepsilon > 0$

$$\psi_\varepsilon(x) := \inf_{y \in K} \left\{ \psi_K(y) + \frac{1}{2\varepsilon} |x - y|^2 \right\}, \quad \forall x \in \mathbb{R}^n.$$

Notice that the inf-convolution of  $\psi_K$  coincides with the Moreau envelope of  $\psi_K$ , denoted by  $e_\varepsilon \psi_K$ : indeed, for all  $x \in \mathbb{R}^n$ ,  $\psi_\varepsilon(x) = \inf_{y \in \mathbb{R}^n} \left\{ \psi_K(y) + \frac{1}{2\varepsilon} |x - y|^2 \right\} =: e_\varepsilon \psi_K(x)$ .

We introduce also the notion of proximal mapping associated to the indicator function  $\psi_K$ , here denoted by  $P_\varepsilon \psi_K$  or simply by  $P_\varepsilon$ :

$$P_\varepsilon(x) = P_\varepsilon \psi_K(x) := \arg \min \left\{ \psi_K(y) + \frac{1}{2\varepsilon} |x - y|^2 \mid y \in \mathbb{R}^n \right\}.$$

For general properties of Moreau envelopes and proximal mappings we refer the reader Rockafellar, Wets (1998).

By  $K_{r_0}$  we denote the set of the points of  $\mathbb{R}^n$ , whose distance from  $K$  is less than  $r_0$ :

$$K_{r_0} := \{x \in \mathbb{R}^n : d_K(x) < r_0\}.$$

Observe that  $P_\varepsilon$  is a single-valued map on  $K_{r_0}$ . Indeed, if  $x \in K$  then obviously  $P_\varepsilon(x) = x$ . So, we assume that  $y_1, y_2 \in P_\varepsilon(x)$  with  $x \in U_K(r_0) = K_{r_0} \setminus K$ . Then, clearly  $y_i \in K$  and  $|x - y_1| = |x - y_2| = d_K(x) < r_0$ . Since  $\Pi_K$  is single-valued on  $U(r_0)$  by Proposition 1, we have that  $y_1 = y_2 = \Pi_K(x)$ . This automatically implies also that if  $x \in K_{r_0}$  then  $\psi_\varepsilon(x) = \frac{1}{2\varepsilon}[d_K(x)]^2 = \frac{1}{2\varepsilon}|x - P_\varepsilon(x)|^2$ .

REMARK 3 Notice that, for all  $x \in K_{r_0}$ ,  $P_\varepsilon(x)$  has the property of being the closest point of  $K$  to  $x$  and it does not depend on  $\varepsilon$ . So, henceforth, in order to simplify the notations we drop the “ $\varepsilon$ ”, writing just  $P(x)$ .

Now, if  $x_1, x_2 \in K_{r_0}$ , then we have

$$\frac{(x_i - P(x_i))}{\varepsilon} \in N_K(P(x_i)) \quad i = 1, 2$$

and by (5)

$$\langle (x_1 - P(x_1)) - (x_2 - P(x_2)), P(x_1) - P(x_2) \rangle + C_0(|x_1 - P(x_1)| - |x_2 - P(x_2)|)|P(x_1) - P(x_2)|^2 \geq 0.$$

Therefore, we obtain

$$|P(x_1) - P(x_2)| \leq \frac{|x_1 - x_2|}{1 - C_0(d_K(x_1) + d_K(x_2))}. \tag{10}$$

Moreover, we get

$$\begin{aligned} \psi_\varepsilon(x_1) - \psi_\varepsilon(x_2) &= \frac{1}{2\varepsilon}(|x_1 - P(x_1)|^2 - |x_2 - P(x_2)|^2) = \\ &= \frac{1}{\varepsilon} \langle x_2 - P(x_2), x_1 - x_2 - (P(x_1) - P(x_2)) \rangle + \frac{1}{2\varepsilon} |x_1 - x_2 - (P(x_1) - P(x_2))|^2 \geq \\ &\geq \frac{1}{\varepsilon} \langle x_2 - P(x_2), x_1 - x_2 \rangle + \frac{1}{\varepsilon} \langle x_2 - P(x_2), P(x_2) - P(x_1) \rangle \geq \\ &\geq \frac{1}{\varepsilon} \langle x_2 - P(x_2), x_1 - x_2 \rangle - \frac{C_0}{\varepsilon} |x_2 - P(x_2)| |P(x_2) - P(x_1)|^2 \end{aligned} \tag{11}$$

where we use (4) since  $x_i - P(x_i) \in N_K(P(x_i))$  and  $P(x_i) \in K$ . So, (2) and (10) yield

$$\psi_\varepsilon(x_1) - \psi_\varepsilon(x_2) \geq \frac{1}{\varepsilon} \langle x_2 - P(x_2), x_1 - x_2 \rangle + O(|x_1 - x_2|^2) \quad \text{as } x_1 \rightarrow x_2.$$

By a similar argument we have

$$\psi_\varepsilon(x_2) - \psi_\varepsilon(x_1) \geq \frac{1}{\varepsilon} \langle x_2 - P(x_2), x_2 - x_1 \rangle + O(|x_1 - x_2|^2) \quad \text{as } x_1 \rightarrow x_2$$

and, so

$$\psi_\varepsilon(x_1) = \psi_\varepsilon(x_2) + \frac{1}{\varepsilon} \langle x_2 - P(x_2), x_1 - x_2 \rangle + O(|x_1 - x_2|^2) \quad \text{as } x_1 \rightarrow x_2.$$

We summarize all the remarks above in the following lemma, whose proof is actually rather standard, but we decided to write it here for the sake of the completeness and also because the introduced arguments will be successively used.

LEMMA 2 *Assume that  $K$  is uniformly prox-regular with constant  $\frac{1}{r}$  for any  $r \in (0, r_0)$ . Then, we obtain the following properties (for any  $\varepsilon > 0$ ):*

1. *the proximal mapping  $P_\varepsilon$  is single-valued on  $K_{r_0}$ ;*
2. *for all  $x \in K_{r_0}$   $\psi_\varepsilon(x) = \frac{1}{2\varepsilon} d_K^2(x)$ ;*
3.  *$\psi_\varepsilon \in \mathcal{C}^{1,1}(K_{r_0})$  and  $\nabla \psi_\varepsilon(x) = \frac{1}{\varepsilon}(x - P_\varepsilon(x)) \in \partial_- \psi_K(P_\varepsilon(x))$  for each  $x \in K_{r_0}$ .*

Next, we prove the existence and uniqueness result of the deterministic Skorokhod problem.

THEOREM 1 *Assume that  $K$  is uniformly prox-regular with constant  $\frac{1}{r}$  for any  $r \in (0, r_0)$  and suppose that (7) holds. Then, for any  $T > 0$  and for any initial data  $x \in K$  the deterministic Skorokhod problem (6) has a unique solution.*

*Proof.* First, we prove the existence. Let us fix  $T > 0$  and chose a point  $x \in K$ . We start solving an approximated Skorokhod problem on  $[0, T]$

$$\begin{cases} \dot{y}_\varepsilon(t) = f(t, y_\varepsilon(t)) - \nabla \psi_\varepsilon(y_\varepsilon(t)) \\ y_\varepsilon(0) = x \in K. \end{cases} \quad (12)$$

By the regularity of  $\nabla \psi_\varepsilon(x)$  (see Lemma 2), the system (12) admits a unique absolute continuous solution  $y_\varepsilon(t)$  as long as  $y_\varepsilon(t) \in K_{r_0}$ . Multiplying by  $\dot{y}_\varepsilon$  the terms of the equation in (12) and integrating on  $[0, t]$  with  $0 < t \leq T$ , we obtain

$$\int_0^t |\dot{y}_\varepsilon(s)|^2 ds + \int_0^t \langle \nabla \psi_\varepsilon(y_\varepsilon(s)), \dot{y}_\varepsilon(s) \rangle ds = \int_0^t \langle f(s, y_\varepsilon(s)), \dot{y}_\varepsilon(s) \rangle ds.$$

Recalling that  $\psi_\varepsilon(x) = 0$  because  $x \in K$ , we obtain

$$\int_0^t |\dot{y}_\varepsilon(s)|^2 ds + \psi_\varepsilon(y_\varepsilon(t)) = \int_0^t \langle f(s, y_\varepsilon(s)), \dot{y}_\varepsilon(s) \rangle ds \leq M\sqrt{T} \left( \int_0^t |\dot{y}_\varepsilon(s)|^2 ds \right)^{\frac{1}{2}}.$$



Hence, we have

$$\int_0^t |\dot{y}_\varepsilon(s)|^2 ds + \psi_\varepsilon(y_\varepsilon(t)) \leq M\sqrt{T} \left( \int_0^t |\dot{y}_\varepsilon(s)|^2 ds \right)^{\frac{1}{2}},$$

that, in turn, yields the following important two estimates that hold true for any  $t \in [0, T]$  as long as  $y_\varepsilon(s) \in K_{r_0}$ :

$$\|\dot{y}_\varepsilon\|_{L^2(0,t)} \leq M\sqrt{T} \tag{13}$$

and

$$\psi_\varepsilon(y_\varepsilon(t)) \leq M^2T. \tag{14}$$

Therefore, as long as  $y_\varepsilon(s) \in K_{r_0}$ , since

$$\psi_\varepsilon(y_\varepsilon(t)) = \frac{1}{2\varepsilon} |y_\varepsilon(t) - P(y_\varepsilon(t))|^2 = \frac{1}{2\varepsilon} d_K^2(y_\varepsilon(t))$$

(see Lemma 2), the inequality (14) implies that

$$d_K(y_\varepsilon(t)) \leq M\sqrt{2T\varepsilon}. \tag{15}$$

Thus,  $y_\varepsilon(t)$  belongs to  $K_{r_0}$  for all  $t \in [0, T]$  whenever

$$0 < \varepsilon < \frac{r_0^2}{2TM^2}. \tag{16}$$

We summarize what we obtained so far: if (16) is satisfied, then system (12) admits a unique solution  $y_\varepsilon$  on  $[0, T]$  and  $y_\varepsilon(t) \in K_{r_0}$  for all  $t \in [0, T]$ . Moreover, thanks to (13) there exists a sequence  $\{\varepsilon_j\}_{j \in \mathbb{N}} \subset \left(0, \frac{r_0^2}{2TM^2}\right)$  converging to zero such that

$$\begin{cases} y_{\varepsilon_j} \rightarrow y \text{ in } \mathcal{C}([0, T]) \\ y_{\varepsilon_j} \rightharpoonup y \text{ in } H^1(0, T) \end{cases}$$

for some  $y \in \mathcal{C}([0, T]) \cap H^1(0, T)$ .

Now, we have to check that  $y$  is really a good candidate, namely

$$\dot{y}(t) \in f(t, y(t)) - \partial_- \psi_K(y(t)) \text{ for a.e. } t \in [0, T].$$

Since

$$f(t, y_\varepsilon(t)) - \dot{y}_\varepsilon(t) = \nabla \psi_\varepsilon(y_\varepsilon(t)) \in \partial_- \psi_K(P(y_\varepsilon(t))),$$

by (4) for all  $z \in K$  we obtain

$$\langle f(t, y_\varepsilon(t)) - \dot{y}_\varepsilon(t), P(y_\varepsilon(t)) - z \rangle + C_0 |f(t, y_\varepsilon(t)) - \dot{y}_\varepsilon(t)| |P(y_\varepsilon(t)) - z|^2 \geq 0$$

and, so, for any  $\varphi \in \mathcal{C}([0, T])$  such that  $\varphi \geq 0$

$$\int_0^T \left\{ \langle f(t, y_\varepsilon(t)) - \dot{y}_\varepsilon(t), P(y_\varepsilon(t)) - z \rangle + C_0 |f(t, y_\varepsilon(t)) - \dot{y}_\varepsilon(t)| |P(y_\varepsilon(t)) - z|^2 \right\} \varphi(t) dt \geq 0.$$

In order to simplify notations, we set  $\xi_\varepsilon(t) := f(t, y_\varepsilon(t)) - \dot{y}_\varepsilon(t)$  and, splitting the above expression into two integrals, we get

$$\int_0^T \langle \xi_\varepsilon(t), P(y_\varepsilon(t)) - z \rangle \varphi(t) dt + C_0 \int_0^T \left\{ |\xi_\varepsilon(t)| |P(y_\varepsilon(t)) - z|^2 \right\} \varphi(t) dt \geq 0.$$

Letting  $\varepsilon \rightarrow 0^+$  (taking a subsequence, if necessary), by (13) the measure  $|\xi_\varepsilon| dt$  weakly converges to some measure  $d\mu(t)$  and we obtain

$$\int_0^T \langle \xi(t), (P(y(t)) - z) \rangle \varphi(t) dt + C_0 \int_0^T \left\{ |P(y_\varepsilon(t)) - z|^2 \right\} \varphi(t) d\mu(t) \geq 0,$$

where  $\xi$  is such that  $\xi_{\varepsilon_j} \rightharpoonup \xi$  in  $L^1(0, T)$ . If we define the measure  $d\lambda(t) := |\xi(t)| dt$ , it turns out that  $d\lambda \ll d\mu$ ; in other words, by the theorem of Radon-Nikodym, there exists a function  $h \in L^1(\mu)$  such that  $d\lambda = h d\mu$ . Finally, the following inequality holds

$$\int_0^T \left\{ \left\langle h(t) \frac{\xi(t)}{|\xi(t)|}, P(y(t)) - z \right\rangle + C_0 |P(y_\varepsilon(t)) - z|^2 \right\} \varphi(t) d\mu(t) \geq 0.$$

Hence, for any countably dense set  $D \subset K$ , we have

$$\left\langle h(t) \frac{\xi(t)}{|\xi(t)|}, P(y(t)) - z \right\rangle + C_0 |P(y(t)) - z|^2 \geq 0 \quad (17)$$

for each  $t \in [0, T] \setminus J_z$  with  $\mu(J_z) = 0$  and  $z \in D$ : the inequality (17) holds true for all  $z$  fixed, so the set of zero measure  $J_z$  could depend on  $z$ ; setting  $Z$  as the union of  $J_z$  such that  $z$  is in a countably dense subset  $D$ , then  $Z$  has zero measure:

$$\mu(Z) = \mu\left(\bigcup_{z \in D} J_z\right) = 0.$$

Inequality (15) implies that  $y(t) \in K$  and, hence,  $P(y(t)) = y(t)$  ( $P(y(t))$  is the closest point of  $K$  to  $y(t)$ ). So, we get

$$\left\langle h(t) \frac{\xi(t)}{|\xi(t)|}, P(y(t)) - z \right\rangle + C_0 |y(t) - z|^2 \geq 0 \quad (18)$$

for all  $t \in [0, T] \setminus Z$  and, moreover, for each  $z \in K$ : indeed, for any  $z \in K$  we can consider a sequence  $z_k \in D$  which converges to  $z$  and such that the above estimate (18) is satisfied. By Remark 1

$$h(t) \frac{\xi(t)}{|\xi(t)|} \in \partial_- \psi_K(y(t)) \quad \text{for a.e. } t \in [0, T]$$

and, hence, we obtain

$$\dot{y}(t) \in f(t, y(t)) - \partial_- \psi_K(y(t)) \quad \text{for a.e. } t \in [0, T].$$

Notice that if  $\dot{y}(t) \in \Pi_{T_K(x(t))}(f(t, y(t)))$  then

$$|\dot{y}(t)| = |\Pi_{T_K(x(t))}(f(t, y(t)))| \leq |f(t, y(t))| < M.$$

So, actually, in order to solve the deterministic Skorokhod problem, it is not necessary to consider the whole  $\partial_- \psi_K(y(t))$  in system (6), but it is sufficient to consider only the set

$$\partial_-^b \psi_K(y(t)) := \{\xi(t) \in \partial_- \psi_K(y(t)) : |\xi(s)| < 2M\},$$

with  $M > 0$  being the constant defined in (7). Indeed, since systems (8) and (9) are equivalent, if  $\xi(t) = f(t, y(t)) - \dot{y}(t) \in \partial_- \psi_K(y(t))$  then  $|\xi(t)| < 2M$ , namely,  $\xi(t) \in \partial_-^b \psi_K(y(t))$ .

Thus, once the initial datum  $x$  is fixed, we found a solution, say  $y_x(\cdot)$ , to the following system with reflection on the boundary

$$\begin{cases} \dot{y}(t) \in f(t, y(t)) - \partial_-^b \psi_K(y(t)) \\ y(t) \in K \quad \forall t \in [0, T] \\ y(0) = x \in K. \end{cases} \tag{19}$$

The uniqueness is an immediate consequence of the following Lemma.

LEMMA 3 *Under the assumptions of Theorem 1, there exists  $\lambda_0 > 0$  such that if  $y_1(\cdot)$  and  $y_2(\cdot)$  are solutions to (19) with initial condition  $y_1(0) = x_1$  and  $y_2(0) = x_2$  respectively, then we have*

$$|y_1(t) - y_2(t)| \leq e^{\lambda_0 t} |x_1 - x_2| \quad \forall t \in [0, T].$$

*Proof.* The proof is an easy consequence of Proposition 1 and Gronwall's inequality. Indeed, let us consider two starting points  $x_1, x_2 \in K$  and two corresponding solution of the Skorokhod problem (19), say  $y_1(\cdot)$  and  $y_2(\cdot)$ . So there exist elements  $\xi_i(t) \in N_K(y_i(t)) \cap \text{Int}(B(0; 2M))$  such that

$$y_i(t) = f(t, y_i(t)) - \xi_i(t) \quad \forall t \in [0, T].$$

By property 4 of Proposition 1 we obtain

$$\frac{r_0}{2M} \langle \xi_1(t) - \xi_2(t), y_1(t) - y_2(t) \rangle \geq -|y_1(t) - y_2(t)|^2.$$

Therefore, we have

$$\begin{aligned} |y_1(t) - y_2(t)| \frac{d}{dt} |y_1(t) - y_2(t)| &\leq \\ &\leq \langle y_1(t) - y_2(t), f(t, y_1(t)) - f(t, y_2(t)) \rangle - \langle \xi_1(t) - \xi_2(t), y_1(t) - y_2(t) \rangle \leq \\ &\leq L|y_1(t) - y_2(t)|^2 + \frac{2M}{r_0} |y_1(t) - y_2(t)|^2 = \left( L + \frac{2M}{r_0} \right) |y_1(t) - y_2(t)|^2 \end{aligned}$$

and, so,

$$\frac{d}{dt} |y_1(t) - y_2(t)| \leq \left( L + \frac{2M}{r_0} \right) |y_1(t) - y_2(t)|.$$

By applying Gronwall's inequality we deduce that

$$|y_1(t) - y_2(t)| \leq e^{\left( L + \frac{2M}{r_0} \right) t} |x_1 - x_2| \quad \forall t \in [0, T].$$

Thus, the statement follows for  $\lambda_0 = L + \frac{2M}{r_0}$ . ■

Now, the proof of Theorem 1 is complete. ■

REMARK 4 *Thanks to the equivalence between (8) and (9) one can study our problem by using an approach in the frame of Viability Theory. Indeed, for all  $(t, y) \in [0, T] \times K$  let us define the function  $\rho(t, y) := d_{T_K(y)}(f(t, y))$  and let us consider the following differential inclusion*

$$\begin{cases} \dot{y}(t) \in B(f(t, y(t)); \rho(t, y(t))) \\ y(t) \in K \quad \forall t \in [0, T] \\ y(0) = x \in K. \end{cases} \quad (20)$$

*Since the set-valued map  $(t, y) \rightsquigarrow B(f(t, y); \rho(t, y))$  turns out to be measurable in  $t$  and upper semicontinuous in  $y$  (see Aubin, Frankowska, 1990), we can apply the Measurable Viability Theorem of Frankowska, Plaskacz (1996), obtaining a viable solution  $y(\cdot)$  to (20) that, in turn, is a solution to (8) and (9). Finally, Lemma 3 guarantees the uniqueness of the solution. Nevertheless, our aim is to show that it is possible to prove Theorem 1 by studying directly the system (6), i.e. without applying existence results for equivalent problems.*

REMARK 5 *It is possible to prove that the above result still holds true when we consider the deterministic Skorokhod problem with oblique reflection:*

$$\begin{cases} \dot{y}(t) \in f(t, y(t)) - A(y(t))\partial_- \psi_K(y(t)) \\ y(0) = x \in K \end{cases}$$

where for each  $x \in \mathbb{R}^n$   $A(x)$  is a symmetric  $n \times n$ -matrix such that  $A : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is of class  $C^1$  with bounded differential and, moreover, there exists  $\nu > 0$  such that

$$\nu I \leq A(x) \leq \nu^{-1} I \quad \forall x \in \mathbb{R}^n.$$

For the details of the proof see Ishii (1988) (and also Appendix B in Bettiol, 2002).

### 3. The Skorokhod control problem and some applications

In this section we deal with the following differential game with reflection on the boundary.

Let us consider the functions  $g : \mathbb{R}^n \times U \times V \rightarrow \mathbb{R}^n$  and  $\ell : \mathbb{R}^n \times U \times V \rightarrow \mathbb{R}$  where  $U$  and  $V$  are compact subsets of  $\mathbb{R}^m$ . We assume that there exists  $M > 0$  such that for all  $x_1, x_2 \in \mathbb{R}^n, u_1, u_2 \in U, v_1, v_2 \in V$

$$\begin{cases} \|g(x, u, v)\|_\infty, \|\ell(x, u, v)\|_\infty \leq M; \\ |g(x_1, u_1, v_1) - g(x_2, u_2, v_2)| \leq M(|x_1 - x_2| + |u_1 - u_2| + |v_1 - v_2|); \\ |\ell(x_1, u_1, v_1) - \ell(x_2, u_2, v_2)| \leq M(|x_1 - x_2| + |u_1 - u_2| + |v_1 - v_2|). \end{cases} \quad (21)$$

Let us denote the set of controls of the first player, called Ursula, and of the second player, called Victor, by  $\mathcal{U} := \{u : [0, +\infty[ \rightarrow U \text{ measurable}\}$  and  $\mathcal{V} := \{v : [0, +\infty[ \rightarrow V \text{ measurable}\}$ , respectively. As usual, we denote an element of  $\mathcal{U}$  by  $u(\cdot)$  or simply by  $u$  when it is understood we are using measurable and bounded functions (and similarly for elements in  $\mathcal{V}$ ).

Once the controls  $u(\cdot) \in \mathcal{U}$  and  $v(\cdot) \in \mathcal{V}$  are fixed, for any  $T > 0$  the following deterministic Skorokhod control problem

$$\begin{cases} \dot{y}(t) \in g(y(t), u(t), v(t)) - \partial_- \psi_K(y(t)) \\ y(t) \in K \quad \forall t \in [0, T]. \\ y(0) = x \in K \end{cases} \quad (22)$$

reduces to (6) simply by taking  $f(t, x) = g(x, u(t), v(t))$ . So, by Theorem 1 for any  $T > 0$ , once we fix  $u(\cdot) \in \mathcal{U}$  and  $v(\cdot) \in \mathcal{V}$ , we get a unique solution  $y_x(\cdot)$  to the differential inclusion (22) starting from the initial data  $x \in K$ .

Take a discount factor  $\lambda > 0$ . In our differential game, Ursula wants to minimize the cost functional

$$\int_0^\infty \ell(y_x(s), u(s), v(s)) e^{-\lambda s} ds,$$

while Victor has to maximize it, where  $y_x(\cdot) \in W_{loc}^{1,1}(0, +\infty)$  is the solution to the controlled Skorokhod problem associated to  $u$  and  $v$ .

For any  $\lambda > 0$ , the (lower) value functions  $w_\lambda$  is so defined:

$$w_\lambda(x) := \inf_{\alpha \in S_U} \sup_{v \in \mathcal{V}} \int_0^\infty \ell(y_x(s), \alpha[v](s), v(s)) e^{-\lambda s} ds \quad \text{for all } x \in K,$$

where  $S_U$  denotes the nonanticipative strategies set of the first player, namely the set of the maps  $\alpha : \mathcal{V} \rightarrow \mathcal{U}$  such that if, for any  $t > 0$  and  $v(\cdot), v'(\cdot) \in \mathcal{V}$ ,  $v(s) = v'(s) \forall s \leq t$  implies  $\alpha[v](s) = \alpha[v'](s) \forall s \leq t$ .

Let us consider the constant  $\lambda_0 > 0$  defined in the proof of Lemma 3. We get the following Hölder estimate on the value function  $w_\lambda$ :

**PROPOSITION 2** *Assume that for some fixed  $r_0 > 0$  the closed set  $K \subset \mathbb{R}^n$  is uniformly prox-regular with constant  $\frac{1}{r}$  for any  $r \in (0, r_0)$  and (21). Take any  $x_0 \in K$  and  $R > 0$ . For all  $\lambda > 0$  such that  $\lambda < \lambda_0$ , we have the following estimate*

$$|\lambda w_\lambda(x_1) - \lambda w_\lambda(x_2)| \leq C |x_1 - x_2|^{\frac{\lambda}{\lambda_0}} \quad \forall x_1, x_2 \in K \cap B(x_0, R) \quad (23)$$

where  $C = \frac{2M}{R+1}$  (it does not depend on  $\lambda$ ).

*Proof.* The proof follows by applying a standard argument based on the Lipschitz dependence of the trajectories on the initial data provided by Lemma 4 (see for instance Bettiol, 2005). ■

It is well known, through the classical theory, that the value function  $w_\lambda$  provides the viscosity solutions of the following stationary Hamilton-Jacobi-Isaacs equation (see Bardi, Capuzzo-Dolcetta, 1997, and Barles, 1994, for the general theory, and Lions, 1985, for the Neumann type boundary conditions):

$$\lambda w_\lambda + H(x, Dw_\lambda) = 0,$$

coupled with the Neumann boundary conditions:

$$\frac{Dw_\lambda}{D\gamma} = 0 \quad \text{on } \partial K,$$

where we suppose that  $K$  is uniformly prox-regular with  $\partial K \in \mathcal{C}^1$  and  $\gamma = \gamma(x)$  is a smooth vector field on  $\partial K$  pointing outwards i.e.

$$\exists \nu > 0 \text{ s. t. } \forall x \in \partial K \quad \langle n(x), \gamma(x) \rangle \geq \nu.$$

The Hamiltonian function  $H$  is given by

$$H(x, p) := \min_{v \in \mathcal{V}} \max_{u \in \mathcal{U}} \{-g(x, u, v) \cdot p - \ell(x, u, v)\}.$$

**REMARK 6** *The Hölder regularity expressed in (23), not only is interesting by itself, but is quite useful for other important properties of the value function.*

Indeed, if, for instance, the domain  $K$  is not only uniformly prox-regular with  $\partial K \in \mathcal{C}^1$  but is also connected and bounded, then we can study the asymptotic behaviour of the term  $\lambda w_\lambda$  as  $\lambda \rightarrow 0^+$ : it is the so-called ergodic problem (for a general introduction to this problem we refer the reader to Bardi, Capuzzo-Dolcetta, 1997). Under a uniform approximate controllability assumption for one player it turns out that there exists a unique  $\chi_0 \in \mathbb{R}$  such that for all  $x \in K$ , we have (see Bettiol, 2005, for the proof):

$$\lim_{\lambda \rightarrow 0^+} \lambda w_\lambda(x) = \chi_0.$$

### Acknowledgments

The author is grateful to both referees for their remarks.

### References

- AUBIN, J.-P. and FRANKOWSKA, H. (1990) Set-valued analysis. *Systems & Control: Foundations & Applications* **2**. Birkhäuser Boston, Basel, Berlin.
- BARDI, M. and CAPUZZO-DOLCETTA, I. (1997) *Optimal control and viscosity solutions of the Hamilton-Jacobi equations*. Birkhäuser, Boston.
- BARLES, G. (1994) *Solutions de viscosité des équations de Hamilton-Jacobi. (French) [Viscosity solutions of Hamilton-Jacobi equations]*. Mathématiques & Applications [Mathematics & Applications], **17**, Springer-Verlag, Paris.
- BETTIOL, P. (2002) Weak Solutions in Hamilton-Jacobi and Control Theory. PhD Thesis.
- BETTIOL, P. (2005) On ergodic problem for Hamilton-Jacobi-Isaacs equations. *ESAIM COCV* **11**, 522–541.
- CLARKE, F.H., STERN, R.J. and WOLENSKI, P.R. (1995) Proximal smoothness and the lower- $\mathcal{C}^2$  property. *J. Convex Analysis* **2**, 117–144.
- CORNET, B. (1983) Existence of slow solutions for a class of differential inclusions. *J. Math. Anal. Appl.* **96**, 130–147.
- DELFOUR, M.C. and ZOLESIO, J.P. (1994) Shape analysis via oriented distance functions. *J. Funct. Anal.* **123** (1), 129–201.
- DELFOUR, M.C. and ZOLESIO, J.P. (2004) Oriented distance function and its evolution equation for initial sets with thin boundary. *SIAM J. Control Optim.* **42** (6), 2286–2304.
- FEDERER, H. (1959) Curvature measures. *Trans. Amer. Math. Soc.* **93**, 418–491.
- FRANKOWSKA, H. (1985) A viability approach to the Skorokhod problem. *Stochastics* **14** (3), 227–244.
- FRANKOWSKA, H. and PLASKACZ, S. (1996) A measurable upper semicontinuous viability theorem for tubes. *Nonlinear Analysis JMA* **26**, 565–582.

- HENRY, C. (1973) An existence theorem for a class of differential equations with multivalued right-hand side. *J. Math Anal. Appl.* **41**, 179–186.
- ISHII, H. (1988) *Lecture Notes on Viscosity Solutions*. Brown University, Providence, RI.
- LIONS, P.L. (1982) *Generalized solutions of Hamilton-Jacobi equations. Research Notes in Mathematics* **69**. Pitman (Advanced Publishing Program), Boston, London.
- LIONS, P.L. (1985) Neumann type boundary conditions for Hamilton-Jacobi equations. *Duke Math. J.* **52** (4), 793–820.
- LIONS, P.L. and SZNITMAN, A.S. (1984) Stochastic differential equations with reflecting boundary conditions. *Comm. Pure Appl. Math.* **37** (4), 511–537.
- ROCKAFELLAR, R.T. and WETS, R.J.-B. (1998) *Variational analysis, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, **317**, Springer-Verlag, Berlin.
- POLIQIN, R.A., ROCKAFELLAR, R.T. and THIBAUT, L. (2000) Local differentiability of distance functions. *Trans. Amer. Math. Soc.* **352** (11), 5231–5249.
- SEREA, O.-S. (2003) On reflecting boundary problem for optimal control. *SIAM J. Control Optim.* **42** (2), 559–575.
- THIBAUT, L. (2003) Sweeping process with regular and nonregular sets. *J. Differential Equation* **193**, 1–26.
- VINTER, R.B. (2000) *Optimal Control*. Birkhäuser, Boston, Basel, Berlin.